

# PRESTOP®

with crops treatments (1/2)

Fungicides			
Active ingredient	Recommended interval (days)	Active ingredient	Recommended interval (days)
Ametoctradin + Dimethomorph	0	Magnesium sulphate + Orange oil (Prev-Magnum)	0
Azoxystrobin	2	Mancozeb	4
Azoxystrobin + Difenconazole	7	Mepanipyrim	0
Benomyl	4	Metalaxyl-M	2
Benthiavalcarb-isopropyl + Folpet	1	Monopotassium phosphonate + Dipotassium phosphonate	2
Bitertanol	2	Myclobutanil	0
Boscalid + Kresoxim-methyl	0	Myclobutanil + Cyclohexanone	0
Bupirimate	0	Myclobutanil + Quinoxifen	1
Carboxin	4	Penconazole	1
Cold pressed orange oil + Hydrocabons aromatic + Alcohol Ethoxylate	2	Potassium bicarbonate	0
		Potassium iodide + Potassium thiocyanate + Addit (Adjuvant)	2**
		Prochloraz	7
Copper oxychloride	0	Procymidone	0
Cyazofamid	1	Propamocarb-HCl	0
Cyflufenamid	0	Propiconazole + Prochloraz	7
Cymoxanil	0	Proquinazid + Tetraconazol	0
Cymoxanil + Zoxamide	1	Pyraclostrobin + Boscalid	2
Difenconazole	7	Pyrimethanil	1
Dithianon	1	Pyriofenone	3
Dodemorph-acetate	0	Quinoxifen	0
Fenhexamid	0	Spiroxamine + Difenconazole	7
Fludioxonil	4	Sulphur	1
Fludioxonil + Cyprodinil	4	Sulphur + Natrium hydroxide	4
Fluopicolide + Propamocarb HCl	1	Tetraconazole	3
Fluopyram	0(1)*	Thiabendazole	4
Fluxapyroxad	0	Tiophanate-methyl	2
Folpet + Dimethomorph	7	Thiram	4
Fosetyl + Propamocarb	0	Toclofos-methyl	2
Fosetyl aluminium	0	Triadimenol	2
Guazatine	2	Trifloxystrobin + Propiconazole	4
Hymexazole	7	Triflumizole	0
Imazalil	2	Vinclozolin	4
Iprodione	4		
Kresoxim-methyl	0		

**0(1)\*** An interval of 1 day is required when PRESTOP® and fluopyram are applied in the same part of the plant. When root system is treated with fluopyram, PRESTOP® can be sprayed in the foliage during the same day.

**2\*\*** PRESTOP® can be sprayed to the foliage two days after application of Enzicur. Re-application of PRESTOP® is necessary after repeated Enzicur treatments.

- ➔ The recommended interval between PRESTOP® and active ingredient or product not mentioned in the list is 7 days.
- ➔ PRESTOP® can be applied in mixtures with compatible products. However, the mixture should be applied immediately after preparation.
- ➔ Do not tank mix PRESTOP® with concentrated solutions of pesticides or fertilizers.

# PRESTOP®

with crops treatments (2/2)

Insecticides		Herbicides	
Active ingredient	Recommended interval (days)	Active ingredient	Recommended interval (days)
Buprofezin	0	2,4 MCPB	4
Chlorpyrifos-methyl	0	Aclonifen	0
Cypermethrin	2	Bentazone	0
Deltamethrine	0	Bentazone + Imazamox	0
Diazinon	0	Ethofumesate	1
Diclorvos	7	Pendimethalin	0
Fenbutatin oxide	2	Prosulfocarb	2
Imidacloprid	0	Pyridate	2
Indoxacarb	0	<b>Microbiological products</b>	
Malathion	0	Active ingredient	Recommended interval (days)
Mevinphos	0	<i>Bacillus amyloliquefaciens</i>	Have some negative effect on the growth of <i>Clonostachys rosea</i> fungus in higher temperature.
Permethrin	2	<i>Bacillus thuringiensis</i>	0
Pirimicarb	0	<i>Beauveria bassiana</i>	0
Pyrethrins	0	<i>Coniothyrium minitans</i>	0
Rape seed oil	0	<i>Lecanicillium muscarium</i>	<i>Clonostachys</i> fungus might have negative effect on the growth of <i>Lecanicillium</i> on the leaf surface.
Spinosad	0	<i>Metarhizium anisopliae</i>	0
Tebufenozide	0	Other microbiological products applied in the growth substrate.	In order to avoid unnecessary competition between different microbes, use the product in rotation, for example at intervals of 7 days.
Growth regulators		Beneficial insects and nematodes	
Product name	Recommended interval (days)	<b>PRESTOP®</b> is compatible with beneficials insects and nematodes.	
Pirouette	0	Disinfectants	
Fertilizers and plant biostimulants		Product name	Recommended interval (days)
Product name	Recommended interval (days)	Resiclean (≤0.01%)	0
IntraCell	0	Huwa-San TR 50 (≤0.04%)	0
Adjuvants		Others	1
Product name	Recommended interval (days)		
Addit	0*		
Break Thru S240	0		
Bluesil SP 3307	0		
Designer	0		
Greenfain	0		
Heliosol	0		
Helioterpen Film	0		
Liberate	1		
Neo-Wett (Imbentin T 080)	0		
Mogiol	0		
PureGreen Spray Oil 13E	0		
Qualibra	0		
Silwet Gold	0		
Sticman	0		
Wetcit	0		

0\* Addit is compatible with PRESTOP® at the maximum concentration of 0,25 %. Higher concentrations of Addit are harmful to PRESTOP®.

- ➔ The recommended interval between PRESTOP® and active ingredient or product not mentioned in the list is 7 days.
- ➔ PRESTOP® can be applied in mixtures with compatible products. However, the mixture should be applied immediately after preparation.
- ➔ Do not tank mix PRESTOP® with concentrated solutions of pesticides or fertilizers.

## LALLEMAND PLANT CARE

LALLEMAND S.A.S. 4, route de Beaupuy – 31180 Castelmorou – France  
 Tel : +33 (0)5 34 27 67 80 – Fax : +33 (0)5 62 79 20 62  
 plantcare@lallemand.com – www.lallemandplantcare.com



LALLEMAND PLANT CARE